

References

- Adam, M., Lacki, J. M. & Barnes, T. G. (1994). Foraging areas and habitat use of the Virginia big eared bat in Kentucky. *J. Wildl. Manage.* 58(3): 462-469.
- Aebischer, N.J., Robertson, P.A. & Kenward, R.E (1993). Compositional analysis of habitat use from animal radio-tracking data. *Ecology* 74(5): 1313-1325.
- Aebischer, N. J. & Robertson, P. A. (1992). Practical aspects of compositional analysis as applied to pheasant habitat utilisation. In *Wildlife Telemetry. Remote Monitoring and Tracking of Animals*: 285-293. Priede, I. G. & Swift, S. M. (eds), Ellis Horwood.
- Amlaner, C.J. & Macdonald, D. (1980). *A Handbook on Biotelemetry and Radio Tracking*. Pergamon Press, Oxford, 804 pp.
- Aldridge H. D. J. N. & Brigham, R. M. (1988). Load carrying and manoeuvrability in an insectivorous bat: A test of the 5 % “rule” of radio-telemetry. *J. Mamm.* 69(2): 379-382.
- Aldridge, H. D. J. N. (1985-86). Manoeuvrability and ecology in British bats. *Myotis* 23-24: 157-160.
- Altmann, J. (1974). Observational study of behavior: sampling methods. *Behaviour* 49: 227-265.
- Arlettaz, R. (1996). Feeding behaviour and foraging strategy of free-living mouse-eared bats, *Myotis myotis* and *Myotis blythii*. *Anim. Behav.* 51(1):1-11.
- Arlettaz, R. (1999). Habitat selection as a major resource partitioning mechanism between the two sympatric sibling bat species *Myotis myotis* and *Myotis blythii*. *Journal of Animal Ecology* 68(3): 460-471.
- Arlettaz, R., Berthoud, G. & M. Desfayes (1998). Tendances démographiques opposées chez deux espèces sympatriques de chauves-souris, *Rhinolophus hipposideros* et *Pipistrellus pipistrellus*: un possible lien de cause à effet?. *Le Rhinolophe* 13: 35-41.
- Arlettaz, R., Godat, S., & Meyer, H. (2000). Competition for food by expanding pipistrelle bat populations (*Pipistrellus pipistrellus*) might contribute to the decline of lesser horseshoe bats (*Rhinolophus hipposideros*). *Biological Conservation* 93(1): 55-60.
- Audet, D. (1990). Foraging behavior and habitat use by a gleaning bat, *Myotis myotis* (Chiroptera: Vespertilionidae). *J. Mamm.* 71: 420-427.
- Barclay, R.M.R. (1989). The effect of reproductive condition on the foraging behavior of female hoary bats, *Lasiurus cinereus*. *Behav. Ecol. Sociobiol.* 24: 31-37.
- Barclay, R.M.R. & Bell, G. (1988). Marking and observational techniques. In: Kunz, T.H.: *Ecological and Behavioral Methods for the Study of Bats*: 59-76.

- Bassi S. & Fabbri I. (1984). Dati preliminari del primo censimento dei Chiroteri delle grotte romagnole. Atti incontro naz. Biospeleologia, Città di Castello: 153-164.
- Baumann, F. (1949). Die freilebenden Säugetiere der Schweiz. Verlag Hans Huber, Bern. 492 pp.
- Beasley, L. J. (1986). Seasonal cycles of pallid bats (*Antrozous pallidus*): proximate factors. *Myotis* 23 – 24: 115–123.
- Beasley, L. J., Pelz, K. M. & Zucker, I. (1984). Circannual rhythms of body weight in pallid bats. *Am. J. Physiol.* 246: 955–958.
- Beck, A., Bontadina, F., Gloor, S., Hotz, T., Lutz, M. & Muehlethaler, E. (1994). Jagdhabitwahl und nächtliche Aufenthaltsgebiete der Grossen Hufeisennase (*Rhinolophus ferrumequinum*) in Raum Castrich/GR. Arbeitsgruppe zum Schutz der Hufeisennasen Graubündens ASGH, Encarden 51, 7152 Sagogn, Switzerland. 102pp.
- Beck, A., Gloor, S., Zahner, M., Bontadina, F., Hotz, T., Lutz, M. & Mühllethaler, E. (1997). Zur Ernährungsbiologie der Grossen Hufeisennase *Rhinolophus ferrumequinum* in einem Alpental der Schweiz. In: (Ohlendorf, B., ed.). Zur Situation der Hufeisennasen in Europa, IFA, Berlin:15-18.
- Beck, A. & Schelbert, B. (1999). Neue Nachweise der Grossen Hufeisennase im Kanton Aargau – Untersuchungen zum Lebensraum und Konsequenzen für den Schutz. *Aarg. Naturf. Ges. Mitt.* 35: 93-113.
- Beck, A., Stutz, H-P. B. & Ziswiler, V. (1989). Das Beutespektrum der Kleinen Hufeisennase *Rhinolophus hipposideros* (Bechstein, 1800) (Mammalia, Chiroptera). *Rev. Suisse Zool.* 96(3): 643-650.
- Begon, M., Harper, J.L. & Townsend, C.R. (1990). Ecology – Individuals, Populations, Communities. Blackwell Scientific Publications, Oxford, GB. 1024 pp.
- Bellamy, P.E., Hinsley, S.A. & Newton, I. (1996). Local extinctions and recolonisations of passerine bird populations in small woods., *Oecologia* 108(1): 64-71.
- Biedermann, M. (1997). Some aspects of the reproduction behaviour of the Lesser Horseshoe Bat (*Rhinolophus hipposideros*) and their consequences for protection. In: (B. Ohlendorf, ed.). Zur Situation der Hufeisennasen in Europa, IFA, Berlin: 77-82.
- Bontadina, F., Arlettaz, R., Fankhauser, T., Lutz, M., Mühllethaler, E., Theiler, A., & Zingg, P. (2000). The lesser horseshoe bat *Rhinolophus hipposideros* in Switzerland: present status and research recommendations. *Le Rhinolophe* 14: 69-83.
- Bontadina, F., Beck, A., Gloor, S., Hotz, T., Lutz, M. & Mühllethaler, E. (1995). Jagt die Grossen Hufeisennase, *Rhinolophus ferrumequinum*, im Wald?

Grundlagen zum Schutz von Jagdgebieten der letzten grössten Kolonie in der Schweiz. Der Ornithologische Beobachter 92: 325-327.

Bontadina, F. & Gloor, S. (1994). How the noctule bat *Nyctalus noctula* selects its roosts in the area of the town of Zurich, Switzerland. Bat Res. News 35(1): 18.

Bontadina, F., Hotz, T., Gloor, S., Beck, A., Lutz, M. & Mühllethaler, E. (1997). Schutz von Jagdgebieten von *Rhinolophus ferrumequinum*. Umsetzung der Ergebnisse einer Telemetrie-Studie in einem Alpental der Schweiz. In: (Ohlendorf, B., ed.). Zur Situation der Hufeisennasen in Europa, IFA, Berlin: 33-39.

Bontadina, F. & Naef-Daenzer, B. (in press). Analysing spatial data of different accuracy: the case of greater horseshoe bats foraging. In: Le Maho, Y.: Wildlife-Biotellemetry, Strassbourg. [online] <http://www.rhinolophus.net/wildtele.pdf>

Bontadina, F., Scaravelli, D., Gloor, S., Hotz, T. & Beck, A. (1999). Radio-tracking bats: a short review with examples of a study in Italy. Atti 1° Convegno Italiano sui Chirotteri, Castell' Azzara: 163–173.

Bontadina, F., H. Schofield & B. Naef-Daenzer (1999). Habitat preference in lesser horseshoe bats as revealed by radio-tracking. Bat Research News 40 (3): 110-111.

Bowman, A.W. (1985). A comparative study of some kernel-based nonparametric density estimators. J. Stat. Comput. and Simulation 21: 313-327.

Breitenmoser, U. (1998). Large predators in the Alps: the fall and rise of man's competitors. Biol. Conserv. 83(3): 279-289.

BUWAL (2000). Waldpolitik. Faktenblätter Bereich Wildtiere. 3.2. Artenvielfalt Fauna. Eidg. Forstdirektion, Bern.

Caccamise, D. F. & Hedin, R. S. (1985). An aerodynamic basis for selecting transmitter loads in birds. Wilson Bull. 97(3): 306–318.

Carroll, C.R. (1992). Ecological management of sensitive natural areas. In: Fiedler, P. L. & Subodh, K. J. (eds). Conservation Biology: the theory and practice of nature conservation, preservation, and management. Chapman and Hall, New York: 347-372.

Catto, C.M.C., Hutson, A.M., Racey, P.A. & Stephenson, P.J. (1996). Foraging behaviour and habitat use of the serotine bat (*Eptesicus serotinus*) in southern England. J. Zool. Lond. 238(4): 623-633.

De Jong, J. (1994). Habitat use, home-range and activity pattern of the northern bat, *Eptesicus nilsoni*, in a hemiboreal coniferous forest. Mammalia 58(4): 535–548.

- De Solla, S. R., Bonduriansky, R. & Brooks, R. J. (1999). Eliminating autocorrelation reduces biological relevance of home range estimates. *J. Anim. Ecol.* 68: 221-234.
- Debinski, D. M. & Holt, R. D. (2000). A survey and overview of habitat fragmentation experiments. *Conserv. Biol.* 14(2): 342-355.
- Dubie, S. & Schwab, F. (1997). Répartition et statut du petit Rhinolophe (*Rhinolophus hipposideros* (Bechstein, 1800) dans le nord et le nord-est de la France. In: (B. Ohlendorf, ed.). *Zur Situation der Hufeisennasen in Europa*, IFA, Berlin: 41-46.
- Duelli, P. (1994). Rote Liste der gefährdeten Tierarten in der Schweiz. BUWAL (Hg.), No. 310.704.d
- Duvergé, L.P. & Jones, G. (1995). Habitat use in greater horseshoe bats; implication for its conservation. In: Gurnell, J: 2nd European Congress on Mammalogy. Abstract Book. SP Press, Cheddar, Somerset, UK.
- Duvergé, P.L (1996). Foraging activity, habitat use, development of juveniles, and diet of the greater horseshoe bat (*Rhinolophus ferrumequinum* - Schreber 1774) in south-west England. Unpublished Ph.D. Thesis. University of Bristol. 310 pp.
- Fairon, J. (1997). Contribution à la connaissance du statut des populations de *Rhinolophus ferrumequinum* et *Rhinolophus hipposideros* en Belgique et problème de leur conservation. In: (B. Ohlendorf, ed.). *Zur Situation der Hufeisennasen in Europa*, IFA, Berlin: 47-54.
- Flückiger, P.F. & Beck. A. (1995). Observations on the habitat use for foraging by *Plecotus austriacus* (Fischer, 1829). *Myotis* 32-33: 121-122.
- Fuhrmann, M. & Seitz, A. (1992). Nocturnal activity of the brown long-eared bat (*Plecotus auritus* L., 1758): data from radio-tracking in the Lenneberg forest near Mainz (Germany). In *Wildlife Telemetry. Remote Monitoring and Tracking of Animals*: 538–548. Priede, I. G. & Swift, S. M. (eds.), Ellis Horwood.
- Gaisler, J. (1963). The ecology of lesser horseshoe bat (*Rhinolophus hipposideros* Bechstein, 1800) in Czechoslovakia, part I. *Vest. Cls. Spol. Zool.* 27: 211–233.
- Gaisler, J. (1989). The r-K selection model and life-history strategies in bats. In *European bat research 1987*: 117-124. Hanak, V., Horacek, I. & Gaisler, J. (eds.), Charles Univ. Press, Praha.
- Gaisler, J. (1997). Preliminary data on the distribution of Rhinolophidae in the Czech Republic and variation in numbers of *R. hipposideros* in S-Moravia. In: (B. Ohlendorf, ed.). *Zur Situation der Hufeisennasen in Europa*, IFA, Berlin: 55-57.
- Geiger, H. (1996). Einsatz der Radiotelemetrie bei Artenschutzbelangen von Fledermaeusen am Beispiel der Grossen Hufeisennase (*Rhinolophus*

ferrumequinum) in Nordbayern. Schriftenreihe für Landschaftspflege und Naturschutz 46: 131-140.

Geiger, H., Hammer, M., Pink, B. & Albrecht, K. (1993). Wochenstabenfund der Grossen Hufeisennase (*Rhinolophus ferrumequinum*) in der Oberpfalz. Koordinationsstelle für Fledermausschutz in Nordbayern, Institut für Zoologie II, Univ. Erlangen. 28 S.

Gellini, S., Casini, L. & Matteucci, C., eds. (1992). Atlante dei Mammiferi della Provincia di Forlì, Maggioli ed.

Grumbine, R. E. (1994). What is ecosystem management? Conserv. Biol. 8: 27-38.

Güttinger, R. (1997). Jagdhabitare des Grossen Mausohrs (*Myotis myotis*) in der modernen Kulturlandschaft. BUWAL-Reihe Umwelt, Nr. 288. Bundesamt für Umwelt, Wald und Landschaft. 140 pp.

Harris, S., Cresswell, W. J., Forde, P. G., Trehewella, W. J., Woollard, T. & Wray, S. (1990). Home-range analysis using radio-tracking data: a review of problems and techniques particularly as applied to the study of mammals. Mamm. Rev. 20: 97-123.

Hickey, M. & Brian, C. (1992). Effects of radiotransmitters on the attack success of hoary bats, *Lasiurus cinereus*. J. Mammal. 73(2): 344-346.

Hooge, P. N. & Eichenlaub, B. (1997). Animal movement extension to ArcView. ver. 1.1. Alaska Biological Science Center, U.S. Geological Survey, Anchorage, AK, USA.

Hughes, P. & Rayner, J. M. V. (1993). The flight of pipistrelle bats *Pipistrellus pipistrellus* during pregnancy and lactation. J. Zool., London 230: 541-555.

Hunter, M. L. (1996). Fundamentals of Conservation Biology. Blackwell Science, Cambridge MA. 482 pp.

Johnson, D. H. (1980). The comparison of usage and availability measurements for evaluating resource preference. Ecology 61(1): 65-71.

Jones, G. & Morton, M. (1992). Radio-tracking studies on habitat use by greater horseshoe bats (*Rhinolophus ferrumequinum*). In Wildlife Telemetry. Remote Monitoring and Tracking of Animals: 521-537. Priede, I. G. & Swift, S. M. (eds). Ellis Horwood.

Jones, G. (1990). Prey selection by the greater horseshoe bat (*Rhinolophus ferrumequinum*): optimal foraging by echolocation? J. Anim. Ecol. 59: 587-602.

Jones, G., Duvergé, P. L. & Ransome, R. D. (1995). Conservation biology of an endangered species: Field studies of greater horseshoe bats. In Ecology, Evolution and Behaviour of Bats. Symposia of the Zoological Society of London 67: 309-324. Racey, P.A. & Swift, S. M. (eds).

- Kenward, R. E. (1992). Quantity versus quality: programmed collection and analysis of radio-tracking data. In *Wildlife Telemetry. Remote Monitoring and Tracking of Animals*: 231–245. Priede, I. G. & Swift, S. M. (eds), Ellis Horwood.
- Kokurewicz, T. (1990). The decrease in abundance of the lesser horseshoe bat *Rhinolophus hipposideros* Bechstein, 1800 (Chiroptera: Rhinolophidae) in winter quarters in Poland. *Myotis* 28: 109–118.
- Kokurewicz, T. (1997). Some aspects of the reproduction behaviour of the Lesser Horseshoe Bat (*Rhinolophus hipposideros*) and their consequences for protection. In *Zur Situation der Hufeisennasen in Europa*: 77–82. Ohlendorf, B. (ed.). Berlin: IFA.
- Krebs, C. J. (1989). Ecological methodology. New York: HarperCollinsPublishers.
- Kronwitter, F. (1988). Population structure, habitat use and activity patterns of the noctule bat, *Nyctalus noctula* Schreb., 1774 (Chiroptera: Vespertilionidae) revealed by radio-tracking. *Myotis* 26: 23–85.
- Krull, D., Schumm, A., Metzner, W. & Neuweiler, G. (1991). Foraging areas and foraging behaviour in the notch-eared bat, *Myotis emarginatus* (Vespertilionidae). *Behav. Ecol. Sociobi.* 28: 247–253.
- Kulzer, E. (1995). Über den Rückgang der Kleinen Hufeisennase *Rhinolophus hipposideros* (Bechstein, 1800) aus Baden-Württemberg. *Laichinger Höhlenfreund* 30: 3–24.
- Kurta, A. & Kunz, T. H. (1987). Size of bats at birth and maternal investment during pregnancy. *Symp. zool. Soc. Lond.* 57: 79–106.
- Lewis, T. & Taylor, L.R. (1964). Diurnal periodicity of flight by insects. *Trans. R. ent. Soc. Lond.* 116(15): 393–476.
- Lugon, A. (1996). Ecologie du Grand rhinolophe *Rhinolophus ferrumequinum* (Chiroptera, Rhinolophidae) en Valais (Suisse): habitat, régime alimentaire et stratégie de chasse. *Trav. Diplome, Univ. Neuchatel.* 116pp.
- Lutz, M. & Muehlethaler, E. (1996). Schutzkonzept fuer die Kleine Hufeisennase *Rhinolophus hipposideros* in den oestlichen Zentralalpen (Lugnez/Valsertal, Graubuenden, Schweiz). In: (Ohlendorf, B., ed.). *Zur Situation der Hufeisennasen in Europa*, IFA, Berlin: 89–94.
- Lyman, C. P. (1970). Thermoregulation and metabolism in bats. pp 301–330. In: *Biology of bats*. Vol. 1. (W. A. Wimsatt, ed). Academic Press, New York. 406 pp.
- MacArthur, R. & Wilson, E.O. (1967). *The Theory of Island Biogeography*. Princeton University Press, Princeton, NJ. 216 pp.
- MacArthur, R.H. & Pianka, E. R. (1966). On optimal use of a patchy environment. *Amer. Natur.* 100: 603–609.

- Mader, H. J. (1984). Animal habitat isolation by roads and agricultural fields. *Biol. Conserv.* 29: 81-96.
- Marzluff, J.M. & Ewing, K. (2001). Restoration of fragmented landscapes for the conservation of birds: a general framework and specific recommendations for urbanizing landscapes. *Rest. Ecol.* 9(3): 280-292.
- McAney, C. M. & Fairley, J. S. (1988). Habitat preference and overnight and seasonal variation in the foraging activity of lesser horseshoe bats. *Acta Theriol.* 33(28): 393–402.
- Mitchell-Jones, A.J. (1995). The status and conservation of horseshoe bats in Britain. *Myotis*, 32/33: 271-284.
- Mitchell-Jones, A.J. (1999). Managing landscapes for the greater horseshoe bat. English Nature, Peterborough. 6 pp.
- Morrissey, W. A., Zinn, J. A. & Corn, M. L. (1994). Ecosystem Management: Federal Agency Activities. Library of Congress, Congressional Research Service, Washington, DC.
- Murphy, D. D., Freas, K. E. & Weiss, S. B. (1990). An environment-metapopulation approach to population viability analysis for a threatened invertebrate. *Conserv. Biol.* 4: 41-51.
- Naef-Daenzer, B. (1993). A new transmitter for small animals and enhanced methods of home-range analysis. *J. Wildl. Manage.* 57(4): 680–689.
- Naef-Daenzer, B. (1994). Grid calculations for homerange and spatial data analysis. Swiss ornithological institute. 41pp.
- Neu, C.W., C.R.Byers, J.M.Peek (1974). A technique for analysis of utilization availability data. *J. Wildl. Manage.* 38(3): 541-545.
- Neuweiler, G. (1989). Foraging ecology and audition in echolocating bats. *Trends Ecol. Evol.* 4(6): 160–166.
- Norberg, U. M. & Rayner, J. M. V. (1987). Ecological morphology and flight in bats (Mammalia; Chiroptera): wing adaptations, flight performance, foraging strategy and echolocation. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 316: 335–427.
- Norusis, M.J. (1986). SPSS/PC + advanced statistics. SPSS Marketing Departement, Chicago.
- Nowak, R. M. (1994). Walker's Bats of the World. Baltimore: Hopkins University Press.
- Ohlendorf, B. (1997a). Zur Situation der Hufeisennasen in Europa, Nebra, 26. - 28. Mai 1995, Tagungsband; Arbeitskreis Fledermaeuse Sachsen-Anhalt e.V., Stecklenberg. IFA Verlag, Berlin. 182 pp.

- Ohlendorf, B. (1997b). Verbreitungsgebiet der Kleinen Hufeisennase (*Rhinolophus hipposideros*) in Europa. In: (Ohlendorf, B., ed.). Zur Situation der Hufeisennasen in Europa, IFA, Berlin: 10-11.
- Otis D. L. & White, G. C. (1999). Autocorrelation of location estimates and the analysis of radiotracking data. *J. Wildl. Manage.* 63(3): 1039–1044.
- Owen, M. (1972). some factors affecting food intake and selection in white-fronted geese. *Journal of Animal Ecology* 41: 79-92.
- Payne, R.J.H. & Wilson, J.D. (1999). Resource limitation in seasonal environments. *OIKOS* Vol. 87(2): 303-314.
- Pfister, M. (2000). Zur Verteilung von Temperatur und Niederschlag im Rhein-/Inngebiet im Hinblick auf eine Schneeschmelz-Abfluss-Simulation. Unpubl. Master Thesis, Univ. Bern, Switzerland.
- Pir, J. (1994). Etho-ökologische Untersuchung einer Wochenstabenkolonie der Grossen Hufeisennase (*Rhinolophus ferrumequinum*, Schreber 1774) in Luxemburg. Unpubl. MSc. thesis, University of Giessen, Germany. 90 pp.
- Priede, I.G. & Swift, S.M., eds (1992). *Wildlife Telemetry: Remote Monitoring and Tracking of Animals*. Ellis Horwood Ltd.. 708pp.
- Primack, R. B. (1998). *Essentials of Conservation Biology*. Sinauer, Sunderland MA.
- Racey, P.A. & Swift, S.M. (1985). Feeding Ecology of *Pipistrellus pipistrellus* (Chiroptera: Vespertilionidae) during Pregnancy and Lactation. I. Foraging Behaviour, II. Diet. *J.anim.Ecol.* 54: 205-215.
- Rackham, O. (1980). *Ancient woodland, its history, vegetation and uses in England*. London: Edward Arnold Publishers Ltd.
- Ransome, R.D. (1968). The distribution of the Greater horse-shoe bat, *Rhinolophus ferrum-equinum*, during hibernation, in relation to environmental factors. *J. Zool.* 154: 77-112.
- Ransome R.D. (1978). Daily activity patterns of the Greater horseshoe bat, *Rhinolophus ferrumequinum*, from April to September. Pp. 259-274 in, R.J. Olembi, J.B. Castelino, F.A. Mutere. (Eds). *Proceedings of the Fourth International Bat Research Conference*. Kenya National Academy for Advancement of Arts and Science, Kenya Literature Bureau, Nairobi.
- Ransome, R.D. (1990). *The natural history of hibernating bats*. Christopher Helm, London. 235 pp.
- Ransome, R.D. & Hutson (2000). Action plan for the conservation of the greater horseshoe bat in Europe (*Rhinolophus ferrumequinum*). Convention on the Conservation of European Wildlife and Natural Habitats Nature and Environment, No. 109. Council of Europe Publishing.

- Rieger, I., Alder, H. & Walzthoeny, D. (1992). Wasserfledermaeuse, *Myotis daubentonii*, im Jagdhabitat ueber dem Rhein. Mitt. Naturf. Gesell. Schaffh. 37: 1-34.
- Rieger I., Walzthoeny, D. & Alder, H. (1990). Wasserfledermäuse, *Myotis daubentonii*, benutzen Flugstrassen. Mitt. Naturf. Gesell. Schaffh. 35: 37-68.
- Robinson, M.F. & Stebbings, R.E. (1997). Home range and habitat use by the serotine bat, *Eptesicus serotinus*, in England. J. Zool. Lond. 243(1): 117-136.
- Roer, H. (1983/84). Zur Bestandessituation von *Rhinolophus ferrumequinum* (Schreber 1774) und *Rhinolophus hipposideros* (Bechstein, 1880) (Chiroptera) im westlichen Mitteleuropa. Myotis 21-22: 122-131.
- Roué, S. (1997). Brève note: Bilan des effectifs observés en 1995 pour les espèces de Rhinolophidae représentées en France. In: (B. Ohlendorf, ed.). Zur Situation der Hufeisennasen in Europa, IFA, Berlin: 133-134.
- Rudolph, B.-U. (1990). Frühere Bestandesdichte und heutige Bestandessituation der Kleinen Hufeisennase *Rhinolophus hipposideros* in Nordbayern. Myotis 28: 101-108.
- Rydell, J. (1992). Exploitation of insects around streetlamps by bats in Sweden. Func. Ecol. 6(6): 744-750.
- Sachs, L. (1992). Angewandte Statistik. Berlin, Heidelberg: Springer-Verlag.
- Sain, S. R., Baggerly, K.A. & Scott, D.W. (1994). Cross-Validation of Multivariate Densities. Journal of American Statistical Association 89: 807-817.
- Saltz, D. (1995). Reporting error measures in radio location by triangulation: a review. J. Wildl. Manage. 58(1): 181-184.
- Saltz, D. & White, G. (1990). Comparison of different measures of the error in simulated radio-telemetry locations. J. Wildl. Manage. 54(1): 169-174.
- Scaravelli, D. & Bassi, S. (1993). Indagini sui chiroteri dell'Appennino Romagnolo-Marchigiano. Biogeographia, 27: 547-552.
- Schnitzler, H. U. & Oswald, J. (1983). Adaptations for the detection of fluttering insects by echolocation in horseshoe bats. In Advances in Vertebrate Neuroethology: 801-827. Ewart, J. P., Capronica, R. R. & Ingle, D. J. (eds.). New York: Plenum press.
- Schober, W. & Grimmberger, E. (1987). Die Fledermäuse Europas, kennen - bestimmen – schützen. Stuttgart, Kosmos Naturführer (Kosmos - Gesellschaft der Naturfreunde). 222 pp.
- Schofield, H. & Morris, C. (1999). The micro-habitat preferences of Bechstein's bat within woodlands in southern England. Bat Res. News 40(3): 140-141.

- Schofield, H. W. (1996). The ecology and conservation biology of *Rhinolophus hipposideros*, the lesser horseshoe bat. PhD thesis, University of Aberdeen.
- Schuler, A. & Bürgi, M. (1998). Wald- und Forstgeschichte. Skript zur Vorlesung 60-316. Proffessur für Forsteinrichtung und Waldwaschstum, Arbeitsbereich Wald- und Forstgeschichte, Zürich: 1-9.
- Schulte, R. (2000). In wieviele Schubladen passt der Naturschutz? - Biotopverbund als integratives Naturschutzkonzept? [online www.nabu-akademie.de/berichte/00_bvs.hmt, 17. 3. 2001]
- Seaman, D. E., Millspaugh, J. J., Kernohan, B. J., Brundige, G. C., Raedeke, K. J. & Gitzen, R. A. (1999). Effects of sample size on kernel home range estimates. *J. Wildl. Manage.* 63(2): 739–747.
- Shaffer, M. (1987). Minimum viable populations: coping with uncertainty. In: Viable populations for conservation (ed: M. E. Soulé), Cambridge University Press, Cambridge: 69-86.
- Sierro, A. & Arlettaz, R. (1997). Barbastelle bats (*Barbastella spp.*) specialize in the predation of moths: Implications for foraging tactics and conservation. *Acta OEcologica*, 18(2): 91-106.
- Silverman, B. W. (1986). Density estimation for statistics and data analysis. Chapman and Hall, London.
- Simberloff, D. (1998). Flagships, umbrellas, and keystones: Is single-species management passé in the landscape era? *Biol. Conserv.* 83(3): 247-257.
- Smith, P. G. (1999). Habitat use by Natterer's bats *Myotis nattereri* on the Welsh/English borders. *Bat Res. News* 40(3): 141-142.
- Smout, T. C. (1997). Scottish woodland history. Edinburgh: Scottish Cultural Press.
- Sokal, R. (1981). Biometry. Freeman & Company, New York, 859pp.
- Soulé, M. E. (1986). Conservation Biology: The Science of Scarcity and Diversity, Sinauer, Sunderland MA.
- Speakman, J.R. & Racey, P.A. (1987). The energetics of pregnancy and lactation in the brown long-eared bat, *Plecotus auritus*. Pp. 367-393 in M.B. Fenton, P.A. Racey, & J.M.V. Rayner (Eds). Recent advances in the study of bats. Cambridge University Press, Cambridge.
- Spitzbergen, F. (1997). Verbreitung und Bestandesentwicklung der Kleinen Hufeisennase (*Rhinolophus hipposideros*) in Österreich. In: (B. Ohlendorf, ed.). Zur Situation der Hufeisennasen in Europa, IFA, Berlin: 135-141.
- Stebbing, R.E. (1982). Radio tracking Greater horseshoe bats with preliminary observations on flight patterns. *Symp. Zool. Soc. London* 49: 161-173.

- Stebbins, R. E. (1988). Conservation of European bats. London: Christopher Helm.
- Stebbins, R.E & Arnold, H.R. (1987). Assessment of trends in size and structure of a colony of the greater horseshoe bat. *Symp. zool. Soc. Lond.* 58: 7-24.
- Stebbins, R.E. & Arnold, H.R. (1989). Preliminary observations of 20th Century changes in distribution and status of *Rhinolophus ferrumequinum* in Britain. In: Hanak, V., I.Horacek & J.Gaisler: European bat research 1987. Charles Univ. Press, Praha: 559-563.
- Stockwell, E. F. (2001). Morphology and flight manoeuvrability in New World leaf-nosed bats (Chiroptera: Phyllostomidae). *J. Zool.*, London 254: 505–514.
- Stutz, H. P. & Haffner, M. (1984). Arealverlust und Bestandesrückgang der Kleinen Hufeisennase (*Rhinolophus hipposideros*) (Bechstein, 1800) (Mammalia: Chiroptera) in der Schweiz. *Jber. Natf. Ges. Graubünden* 101: 169–178.
- Suter, W. (1998). Involving Conservation Biology in biodiversity strategy and action planning. *Biol. Conserv.* 83 (3): 235-237.
- Tanner, K.M. & Zoller, S. (1996). Wie Meliorationen die Ausstattung der Kulturlandschaft verändern. *Info. blatt Forsch. bereich Landschaftsoekologie* 32: 3-6.
- Tufto, J., Andersen, R. & Linnell, J. (1996). Habitat use and ecological correlates of home range size in a small cervid: the roe deer. *Journal of Animal Ecology* 65(6): 715-724.
- Vickery, J.A., Tallowin, J.R., Feber, R.E., Asteraki, E.J., Atkinson, P.W., Fuller, R.J. & Brown, V.K. (2001). The management of lowland neutral grasslands in Britain: effects of agricultural practices on birds and their food resources. *Journal of Applied Ecology* 38(3): 647-664.
- Von der Emde, G. & Menne, D. (1989). Discrimination of insect wingbeat-frequencies by the bat *Rhinolophus ferrumequinum*. *J. Comp. Physiol. A.* 164: 663–671.
- Voûte, A. M., Sluiter, J. W. & van Heerdt, P. F. (1980). De vleemuizenstand in enige Zuidlimburgse groeven sedert 1942. *Lutra* 22(1–3): 18–34.
- Warren R. D. (1998). Lesser horseshoe bat summer roost monitoring, 29 may to 17 June 1998. Countryside Council for Wales Natural Science Report 98/9/2.
- White, G.C. & Garrott, R.A. (1990). Analysis of wildlife radio-tracking data. Academic press, Inc., San Diego, California. 383 pp.
- Wiens J. A. & Stenseth, N. C. (1993). Ecological mechanisms and landscape ecology. *Oikos* 66(3): 369-380.

- Wiens, J.A. & Milne, B.T. (1989). Scaling of “landscapes” in landscape ecology, or, landscape ecology from a beetle’s perspective. *Landscape Ecology* 3: 87-96.
- Wilcox, B. A. (1980). Insular ecology and conservation. In: Soulé, M. E. & Wilcox, B. A.: *Conservation Biology. An evolutionary-ecological perspective*. Sinauer, Sunderland MA.
- Wilkinson, G.S. & Bradbury, J.W. (1988). Radiotelemetrie: techniques and analysis. In: Kunz, T.H.: *Ecological and Behaviour Methods for the Study of bats*. Smiths-Inst. Press, Washington D.C. and London: 105-124.
- Worton, B. J. (1989). Kernel methods for estimating the utilisation distribution in home range studies. *Ecology*. 70: 164–168.
- Worton, B. J. (1987). A review of models of home range for animal movement. *Ecological Modelling* 38: 277-298.
- Worton, B.J. (1995). Using Monte Carlo simulation to evaluate kernel-based home range estimators. *J. Wildl. Manage.* 59(4): 794-800.
- Wray, S., Cresswell, W.J., White, P.C.L. & Harris, S. (1992). What, if anything, is a core area? An analysis of the problems of describing internal range configurations. In: Priede, I.G. & S.M. Swift. *Wildlife Telemetry. Remote Monitoring and Tracking of Animals*. Ellis Horwood, Chichester: 521-537.
- Zahner, M. (1984). Nahrungzusammensetzung, aktivität und nächtliche aufenthaltsgebiete der Grossen Hufeisennase *Rhinolophus ferrumequinum* (Chiroptera: Rhinolophidae). Unpublished PhD. thesis. Switzerland: University of Zurich. 39 pp.
- Zahner, M. (1996). Aktivität und nächtliche Aufenthaltsgebiete der Grossen Hufeisennase *Rhinolophus ferrumequinum* (Chiroptera, Rhinolophidae) in Castrisch (Vorderrheintal, Graubünden). *Jber. Naturf. Ges. Graubünden* 108: 155-173.
- Zimmermann, J. W. & Powell, R. A. (1995). Radiotelemetry error: location error method compared with error polygons and confidence ellipses. *Can. J. Zool.* 73: 1123–1133.